

Original Article

Functional Age and Bouts of Physical Activity in Middle-Aged to Older Japanese Adults; Yurin-StudyMakoto Ayabe¹⁾, Yoshikazu Yonei²⁾, Ryo Miyazaki²⁾, Hideaki Kumahara³⁾, Naoki Sakane¹⁾, Kojiro Ishii⁴⁾

1) Division of Preventive Medicine, Clinical Research Institute, National Hospital Organization Kyoto Medical Center

2) Anti-aging Medical Research Center, Graduate School of Life and Medical Sciences, Doshisha University

3) Faculty of Nutrition Science, Nakamura University

4) Faculty of health & Sports Science, Doshisha University

Abstract**Purpose:** To examine the relationship between the functional age (FA) and the objectively measured daily physical activity (PA).**Methods:** The subjects of the present investigation were 15 males and 22 females (68±6 yr.). The Anti-Aging-Dock determined the 5 components of FA (Bone age; BA, Hormone-age; HA, Muscle-age; MA, Nervous system-age; NA, Vascular-age; VA). Each participant wore an electronic pedometer (HJ-720IT, Omron Corporation, Kyoto-city, Kyoto Japan) in order to determine daily number of steps (STEP_{day}) and the time and steps for PA lasting longer >10-min at >60 steps·min⁻¹ of the step rate (Bout_{time} and Bout_{step}). The HA was evaluated by using the standard aging curve of serum DHEA-s and IGF-I concentration.**Result:** Simple correlation coefficient showed that the HA significantly associated with STEP_{day} and Bout_{step} in female subjects ($p < 0.05$). A multiple regression analysis showed that the Bout_{step} and Age came out significant parameter of the HA.**Conclusions:** The HA significantly associated with steps Bout_{step} independently from the STEP_{day} in middle-aged to older female adults. The bouts of PA lasting >10-min may be an effective PA to protect the age-associated changes in hormonal functions.**KEY WORDS:** Aging, Pedometer, Accelerometer, Exercise, Public health**Introduction**

An active lifestyle is an important protect factor for chronic disease morbidity and mortality in aging¹⁾. One of the main determinants of the levels of physical activity (PA) is age²⁾, and the PA frequently decreased with increasing age³⁾. However, older individuals can still obtain a desirable amount of PA as a result of an appropriate intervention⁴⁾. Furthermore, a prospective cohort study suggested that elderly women could lengthen their life by increasing and maintaining PA levels⁵⁾. Therefore, older individuals should improve their PA level as well as younger- and middle-aged adults in order to elicit positive health benefits.

A consensus statement⁶⁾ recommended that older adults need moderate-intensity aerobic PA for a minimum of 30 min on five days each week or vigorous intensity aerobic PA for a minimum of 20 min on three days each week in order to promote and maintain health. Furthermore, for older adults who are not active at recommended levels, plans should include a gradual approach to increase PA over time using multiple bouts of PA (>10 min)⁶⁾.

The functional age (FA), determined by the Anti-Aging-Dock, is a useful tool for diagnosing the degree of aging⁷⁾. The FA is consisted of five parameters in regard to the psychological and physiological indicator, such as the Bone-age (BA), Hormone-age (HA), Muscular-age (MA), Neurological-age

(NA), and, Vascular-age (VA). Although previous studies suggested that the exercise program could improve some of the components of FA⁸⁻¹²⁾, the changes of the FA after the interventions were inconsistent. These studies mainly used the pedometer-based intervention, thus the exercise prescription had limited to advice the increases in the total amounts of the walking activities. The exercise intensity and the duration are the important determinants of the effects of the exercise intervention^{4,13)}. Therefore, the walking speed was associated with survival in older adults¹⁴⁾. Researchers have also found evidence that older individuals who had participated in the moderate intensity PA on most days of week had a better physical function than older adults who were active throughout the day¹⁵⁾. As previously mentioned, the bouts of PA, such as PA lasting at least 10 minutes, have also been recommended in order to obtain the health benefits, rather than the shorter PA⁶⁾. In the best of our knowledge, the relationships between the FA and the bouts of PA remain unclear. Based on these findings, we hypothesized that the bouts of PA will have a greater effect on FA than the total amount of PA.

Thus, the primary objective of the present investigation was to examine the relationship between each of 5 components of FA and the objectively measured daily PA such as the daily number of steps (STEP_{day}) and the bout of PA lasting longer than 10 minutes.

Methods

Subjects

Fifteen men and twenty-two women, aged from 56 yr. to 79 yr., participated in the present investigation. The characteristics of subjects were shown in [Table 1](#). Subjects were recruited from the Senior Club of Yurin School District, a school district in the Shimogyo district of Kyoto-city, Japan, using bulletin boards in the district. The former Yurin Elementary School (Former-Yurin Sho) is used as a community center. All participants were independently living, and were free from any severe chronic diseases. After explaining the study design and requirements, all subjects signed a consent form. All procedure of the present study was approved by the ethics committee of the Doshisha University.

Physical Activity Assessment

All subjects wore an electronic pedometer (HJ-720IT, Omron Corporation, Kyoto-city, Kyoto Japan) on their waist level just above the leg continuously for 14 days, except while sleeping and bathing. At the end of the data collection period, the device was retrieved, and the data were downloaded into a computer. This type of pedometer is a reliable method of recording the number of steps under controlled and free-living conditions ¹⁶⁻¹⁹. In the present investigation, the daily time (**min·day⁻¹**) and the number of steps (**steps·day⁻¹**) for bout of PA (Bout_{time} and Bout_{step}) were assessed in addition to the STEP_{day}. The bout of PA was defined as the consecutive PA lasting longer than 10 minutes at least 60 steps·min⁻¹ of the minute-by-minute stepping rate. However, it must be noted that, the bouts of PA allowed including the interruptions. Limited to one minute, the minute-by-minute stepping rate might drop below 60 **step·min⁻¹** during a bout.

Functional Age

The FA was determined following the previously described procedures ^{7,9}. The MA was determined based on the muscle mass determined by a bioelectrical impedance analyzer (Physion MD, NIPPON SHOOTER Ltd, Chiyoda-ku, Tokyo, Japan). The HA was evaluated by using the standard aging curve of the serum dehydro-epiandrosterone sulfat (DHEA-s) and insulin-like growth factor-1 (IGF-1). The NA was evaluated according to the results of the Wisconsin card-sorting test. The VA was determined based on photo-plethysmogram (Dynapulse SDP-100, Fukuda Denshi Co., Ltd., Bunkyo-ku Tokyo, Japan). Finally, the BA was determined by the bone mineral density assessed by ultrasound bone densitometer (A-1000, GE Yokogawa Medical Systems Ltd, Hino-city, Tokyo, Japan). Further details can be found in the previous publications ^{7,9}.

Date Analisis and Statistics

Linear regressions analyses were performed for either the FA and the several index of PA, including the bout of PA and the daily total amount of PA, to investigate the ability and contribution for the FA. In addition, a multiple stepwise regression analysis was used to investigate the contribution of the FA to the index of PA and characteristics of subjects. Pearson's correlation coefficients (*r*) as well as product correlation coefficient (*r*²) were calculated. The difference between men and women was analyzed using the Mann-Whitney-U test. All statistical analyses were performed by using the StatView software program (version 5.0.1, SAS Institute, Cary, NC, USA). The data are expressed as the mean with standard deviation. Statistical significance was set at *p* < 0.05.

Results

The characteristics of the subjects are presented in [Table 1](#). The height and body weight were significantly higher in men compared with that in women (*p* < 0.05). The number of steps and the time for the bout of PA are shown in [Table 2](#). In regard to the indexes of PA, although marginal difference was found in STEP_{day} (*p* = 0.07), All variables did not differ significantly between men and women. The BA was significantly younger in male subjects compared with that in female subjects (*p* < 0.05, [Table 3](#)). The MA, HA, NA, and VA did not differ significantly between men and women.

Table 1 Characteristics of subjects in Yurin-study

	All	Women	Men
Age (yr.)	68.0 ± 8.0	68.0 ± 6.0	67.0 ± 6.0
Height (cm)	159.3 ± 7.3	155.1 ± 5.0	165.3 ± 5.7 **
Body weight (kg)	58.6 ± 10.0	54.9 ± 8.2	64.1 ± 10.1 **
Body mass index (kg·m ⁻²)	23.0 ± 2.8	22.8 ± 2.8	23.4 ± 2.7
Waist circumfluence (cm)	85.0 ± 9.0	86.0 ± 9.0	85.0 ± 10.0

Data are expressed as mean with standard deviation (mean±SD).

Significant difference compared with that in women (*p* < 0.01).

Table 2 Number of steps and time for bout and non-bout of physical activity in Yurin-study

	All	Women	Men
Daily number of steps (steps·day ⁻¹)	8691 ± 3822	7811 ± 1886	10072 ± 5479
Bout _{step} (steps·day ⁻¹)	3114 ± 73611	2508 ± 1514	4067 ± 5459
Bout _{time} (min·day ⁻¹)	45 ± 55	45 ± 58	44 ± 53

Data are expressed as mean with standard deviation (mean±SD).

There are no significant difference between men and women.

Bout_{step}; The number of steps for the bout of physical activity,

Bout_{time}; The time for the bout of physical activity.

Table 3 Functional age in Yurin-study

	All	Women	Men
Muscular age (yr.)	55 ± 7	56 ± 7	56 ± 7
Bone age (yr.)	59 ± 14	64 ± 13	53 ± 13*
Hormone age (yr.)	70 ± 8	71 ± 8	68 ± 7
Neurological age (yr.)	63 ± 15	62 ± 15	65 ± 16
Vascular age (yr.)	65 ± 9	65 ± 8	65 ± 9

Data are expressed as mean with standard deviation (mean±SD).

*Significant difference compared with that in women (**p* < 0.05).

There were no significant relationship between the FA and the indexes of PA in men. In contrast, the HA significantly associated with the STEP_{day} and the bout_{step} in female subjects (*p* < 0.05, [Fig. 1](#)). Furthermore, the marginal significant relationships were found between STEP_{day} versus the BA and the bout_{step} versus the MA (*p* = 0.06 and *p* = 0.09).

[Table 4](#) shows the results of the multiple stepwise regression analysis for the HA and PA in female subjects ([Table 4](#)). Because the significant relationship did not found in male subjects, the multiple stepwise regression analysis was limited to the female subjects. In addition to the several indexes of PA (STEP_{day}, Bout_{step} Bout_{time}) and age were included to the independent variables, because age significantly associated with the HA (*p* < 0.05). As the regard to the index of the PA, all indexes




Fig. 1. Relationships between hormonal age and bout and non-bout of physical activity in Yurin-study. Bout_{step}; The number of steps for the bout of physical activity, Bout_{time}; The time for the bout of physical activity.

(STEP_{day}, Bout_{step}, Bout_{time}) were included to the independent variable. The Bout_{step} accounting for 22.4% of the total variance (Table 4). The total amount of age also account for 13% of the variance in the HA, respectively.

Table 4 Multiple stepwise regression analyses for hormone age and daily physical activity in female subjects in Yurin-study

Dependent variable	Medaled variable ^a	SEE	R ²
Hormone age	Step 0. Bout _{step}	7.3	0.224*
	Step 1. Bout _{step} + Age	6.5	0.354**

* $p < 0.05$, ** $p < 0.001$

Bout_{step}; The number of steps for the bout of physical activity, SEE; standard error of mean.

^a Calculated Variables: The analysis employed 4 variables: age, the daily number of steps, the number of steps for the bout of physical activity, the time for the bout of physical activity.

Discussion

The present investigation examined the relationships between the FA and the bout or nonbout of PA in middle-aged to older adults. As results, the HA significantly associated with STEP_{day} and the Bout_{step} in female subjects. Thereafter, the multiple regression analysis showed that the Bout_{step}, rather than STEP_{day}, was a significant contributor of the HA. These results indicate that the higher number of the Bout_{step} associated with the younger HA independent from the STEP_{day} in middle-aged to older female adults. However, the relationships between the indexes of PA and the FA except for the HA remains unclear in particular male adults. These finding suggested that the middle-aged to older women should be encouraged to participate in the PA lasting longer than 10 min at the stepping rate of >60 steps·min⁻¹ in order to improve the hormonal distribution including the IGF-1. Although the significant relationship with Bout_{step} was limited to the HA, the Bouts of PA may elicit several health benefits, because IGF-1 is an important metabolic biomarker associated with a variety of health- and exercise-related outcomes²¹.

The original aspects of the present investigation were to demonstrate the relationships between the FA with the bout and non-bout of PA. In particular, the bout of PA significantly associated with the HA independently from STEP_{day} in female subjects. These results of the present investigation was supported by the previous findings^{9,12}. As results of the pedometer-based walking program, the FA, including the HA, did not differ significantly by increasing STEP_{day} in middle-aged to older adults^{9,12}. In contrast, Mochizuki *et al.*¹¹ showed that the standardized exercise program (40 min·time⁻¹, 3 times·wk⁻¹) could successfully improved the components of the HA such as IGF-1 and DHEA-s. These findings apparently indicate that the accumulating the continuous bout of PA may be preferable strategy to obtain the younger HA rather than the increases in STEP_{day} by taking the intermittent PA.

In regard to the relationship between the HA and the Bout_{step}, although the present investigation cannot explain the physiological mechanism, these results might depend on the IGF-1 responses to PA. Because, the HA was evaluated based on the IGF-1 and DHEA-s, and the additional analysis showed that the IGF-1 significantly associated with the Bout_{step} (Fig. 2). Therefore, the relationships between the HA and the Bout_{step} may depends on the IGF-1 rather than DHEA-s. Several experimental studies showed that the serum IGF-1 response to




Fig. 2. Relationships between insulin-like growth factor 1 age and bout and non-bout of physical activity in Yurin-study. Bout_{step}; The number of steps for the bout of physical activity, Bout_{time}; The time for the bout of physical activity.

the aerobic exercise depended on the intensity and duration^{22,23}). The brief exercise leads to small but significant increases in circulating IGF-I, and these responses may be influenced by exercise intensity²²). Furthermore, the moderate intensity PA of long duration may modulate the bioavailability of IGF-1²³). In the present investigation, the bout was defined as the PA lasting longer than 10-min at 60 steps·min⁻¹ of stepping rate. The stepping rate is a practical indicator of the exercise intensity¹⁷), and the PA at 60 steps·min⁻¹ or higher would be classified as the locomotive activity such as walking and running. In contrast, the PA at <60 steps·min⁻¹ is corresponded to the lifestyle activity such as the housekeeping, dinning, gardening etc²⁴). Furthermore, the duration of PA was mainly less than <30-sec, and the frequency of the PA lasting >10-min was very rare under the free-living condition²⁵). Based on these findings, the Bout_{step} is considered as “a spontaneous exercise” rather than “the usual activity of daily living”. Therefore, it is reasonable to assume that the serum IGF-1 closely associate with the Bout_{step} rather than STEP_{day}.

There are some limitations in regard to the present investigation. First, the number of participant used in the present investigation was relatively small, and were limited to mainly healthy subjects. Second, due to the potential limitation with this device, the Bout_{step} (60 steps·min⁻¹, >10 min) did not completely reflect the intensity and the bouts duration of PA. Not only stepping rate but also stride length contributes to the exercise intensity, and the Bout_{step} allows including the interruptions. Furthermore, the Omron pedometer could be a useful device for the continuous bouts of moderate-intensity walking, but it does not measure all steps taken throughout the day due to the underestimation for very light intensity PA¹⁹). Third, the Bout_{step} and age are account for only 35% of the variations in the HA. The additional determinants of the HA, such as the nutritional and genetic factors, may be examined in the future studies. Finally, the type and position of the pedometer used in the present investigation does not allow for collection of upper body activity and thus may underestimate total PA.

In summary, the present investigation examined the relationships between the FA and the bout and non-bout of PA in middle-aged to older adults. As results, the higher number of the Bout_{step} associated with the younger HA independent from STEP_{day} in middle-aged to older female adults. These findings suggest that the middle-aged to older women should be encouraged to participate in the PA lasting longer than 10 min in order to improve the hormonal distribution including the

IGF-1. The relationships between the indexes of PA and the FA except for the HA remains unclear in particular male adults. Future study should assess the effects of the bout of PA on the several health outcomes in a larger heterogeneous sample.

References

- 1) DiPietro L: Physical activity in aging. *J Gerontol A Biol Sci Med Sci.* 56;13-22: 2001
- 2) Westerterp KR, Meijer EP: Physical Activity and Parameters of Aging. *J Gerontol A Biol Sci Med Sci.* 56;7-12: 2001
- 3) Ayabe M, Yahiro T, Yoshioka M, et al: Objectively measured age-related changes in the intensity distribution of daily physical activity in adults. *J Phys Act Health.* 6;419-25: 2009
- 4) Chodzko-Zajko WJ, Proctor DN, Fiatarone Singh MA, et al: American College of Sports Medicine position stand. Exercise and physical activity for older adults. *Med Sci Sports Exerc.* 41;1510-30: 2009
- 5) Gregg EW, Cauley JA, Stone K, et al: Relationship of changes in physical activity and mortality among older women. *JAMA.* 289;2379-86: 2003
- 6) Nelson ME, Rejeski WJ, Blair SN, et al: Physical activity and public health in older adults: recommendation from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc.* 39;1435-45: 2007
- 7) Yonei Y, Mizuno Y: The human dock of tomorrow: annual health check-up for anti-aging. *Ningen Dock.* 19;5-8: 2005
- 8) Miyazaki R, Hasegawa T, Fujioka N, et al: Effects on anti-aging indicators in middle-aged men of an intervention to prevent lifestyle-related diseases; Pilot study utilizing a company-wide anti-aging medical checkup and pedometers. *Anti-Aging Medicine.* 6;83-94: 2009
- 9) Miyazaki R, Ishii K, Ichikawa H, et al: Community medicine and anti-aging: effects of combining a long-term pedometer-based physical activity program with anti-aging medical checkups on health and anti-aging medical indicators in community-dwelling older adults (yurin study 1). *Anti-Aging Medicine.* 7;143-52: 2010
- 10) Miyazaki R, Yonei Y, Azuma Y, et al: Relationship between the change in daily step count and brachial-ankle wave velocity during a pedometer-based physical activity program for older adults. *Anti-Aging Medicine.* 8;35-40: 2011
- 11) Mochizuki T, Amenomori Y, Miyazaki R, et al: Evaluation of exercise programs at a fitness club in female exercise beginners using anti-aging medical indicators. *Anti-Aging Medicine.* 6;66-78: 2009
- 12) Nomoto K, Miyazaki R, Hasegawa T, et al: Efficacy of a health promotion program with anti-aging medical checkup and instructions for walking under pedometer management in factory workers. *Anti-Aging Medicine.* 7;73-84: 2010
- 13) Yonei Y, Takahashi Y, Hibino S, et al: The effects of walking with pedometers on quality of life and various symptoms and issues relating to aging. *Anti-Aging Medicine.* 5;22-9: 2008
- 14) Studenski S, Perera S, Patel K, et al: Gait speed and survival in older adults. *JAMA.* 305;50-8: 2011.
- 15) Brach JS, Simonsick EM, Kritchevsky S, et al: The association between physical function and lifestyle activity and exercise in the health, aging and body composition study. *J Am Geriatr Soc.* 52;502-9: 2004
- 16) Ayabe M, Aoki J, Ishii K, et al: Pedometer accuracy during stair climbing and bench stepping exercises. *J Sports Sci & Med.* 7;249-54: 2008
- 17) Ayabe M, Kumahara H, Aoki J, et al: Quantifying time spent in moderate to vigorous intensity physical activity via stepping rate. *Jap J Phys Fitness and Sports Med.* 57;453-62: 2008
- 18) Holbrook EA, Barreira TV, Kang M: Validity and reliability of Omron pedometers for prescribed and self-paced walking. *Med Sci Sports Exerc.* 41;670-4: 2009
- 19) Silcott NA, Bassett DR, Jr., Thompson DL, et al: Evaluation of the Omron HJ-720ITC pedometer under free-living conditions. *Med Sci Sports Exerc.* In Press
- 20) Ohshima H: Relationship between walking activity and percent body fat on weekdays and weekend days in female adults *Journal of Japan Society for the Study of Obesity.* 12;119-23: 2006
- 21) Nindl BC, Pierce JR: Insulin-like growth factor I as a biomarker of health, fitness, and training status. *Med Sci Sports Exerc.* 42;39-49: 2010
- 22) Schwarz AJ, Brasel JA, Hintz RL, et al: Acute effect of brief low- and high-intensity exercise on circulating insulin-like growth factor (IGF) I, II, and IGF-binding protein-3 and its proteolysis in young healthy men. *J Clin Endocrinol Metab.* 81;3492-7: 1996
- 23) Chadan SG, Dill RP, Vanderhoek K, et al: Influence of physical activity on plasma insulin-like growth factor-1 and insulin-like growth factor binding proteins in healthy older women. *Mech Ageing Dev.* 109;21-34: 1999
- 24) Ayabe M, Aoki J, Kumahara H, et al: Minute-by-minute stepping rate of daily physical activity in normal and overweight/obese adults. *Obesity Research & Clinical Practice.* 5;151-6: 2011
- 25) Ayabe M, Kumahara H, Aoki J, et al: Assessment of frequency and bouts duration of moderate intensity physical activity by an accelerometer in 4-sec interval under-free living condition. *Journal of Japan Society for the Study of Obesity.* 13;197-200: 2007